

Authors are encouraged to submit new papers to INFORMS journals by means of a style file template, which includes the journal title. However, use of a template does not certify that the paper has been accepted for publication in the named journal. INFORMS journal templates are for the exclusive purpose of submitting to an INFORMS journal and should not be used to distribute the papers in print or online or to submit the papers to another publication.

Random Knockout Tournaments

Ilan Adler

Department of Industrial Engineering and Operations Research, University of California, Berkeley, adler@ieor.berkeley.edu

Yang Cao

Department of Industrial and Systems Engineering, University of Southern California, cao573@usc.edu

Richard Karp

Department of Electrical Engineering and Computer Sciences, University of California, Berkeley, karp@cs.berkeley.edu

Erol A. Peköz

Questrom School of Business, Boston University, pekoz@bu.edu

Sheldon M. Ross

Department of Industrial and Systems Engineering, University of Southern California, smross@usc.edu

We consider a random knockout tournament among players $1, \dots, n$, in which each match involves two players. The match format is specified by the number of matches played in each round, where the constitution of the matches in a round is random. Supposing that there are numbers v_1, \dots, v_n such that a match between i and j will be won by i with probability $v_i/(v_i + v_j)$, we obtain a lower bound on the tournament win probability for the best player, as well as upper and lower bounds for all the players. We also obtain additional bounds by considering the best and worst formats for Player 1 in the special case $v_1 > v_2 = v_3 = \dots = v_n$.

History: This paper was first submitted on December 6, 2016.

1. Introduction

We consider a tournament among players $1, \dots, n$, in which each match involves two players. The tournament is assumed to be of knockout type in that the losers of matches are eliminated and do not move on to the next round, and the tournament continues until all but one player is eliminated, with that player being declared the winner of the tournament. The match format is specified by the set of positive integers r, m_1, \dots, m_r with the interpretation that there are a total of r rounds, with

round i consisting of m_i matches, $\sum_{i=1}^r m_i = n - 1$. Because $\sum_{j=1}^{i-1} m_j$ players have been eliminated by the end of round $i - 1$, we must have that $m_i \leq (n - \sum_{j=1}^{i-1} m_j)/2$.

We suppose that the constitution of the matches in a round is totally random. That is, for instance, the $2m_1$ players that play in round 1 are randomly chosen from all n players and then randomly arranged into m_1 match pairs. The winners of these m_1 matches, along with the $n - 2m_1$ players that did not play a match in round 1 then move to round 2, and so on.

We suppose that the players have respective values v_1, \dots, v_n , and that a match involving players i and j is won by Player i with probability $v_i/(v_i + v_j)$. We let P_i be the probability that Player i wins the tournament, $i = 1, \dots, n$. In Section 3 we derive a lower bound on the probability that the strongest player (e.g., the one with the largest v) wins the tournament, and an upper bound on the probability that the weakest player wins the tournament. In Section 4 we derive upper and lower bounds on the win probabilities $P_i, i \geq 1$, and also show that if $v_1 \geq v_2 \geq \dots \geq v_n$ then $P_1 \geq P_2 \geq \dots \geq P_n$. In Section 5, we consider the special case where $v_1 > v_2 = \dots = v_n$ and show that when $n = 2^s + k, 0 \leq k < 2^s$, the best format for the strongest player is the so-called *balanced format* that has k matches in the first round and then has all remaining players competing in each subsequent round. We also show that whenever the number of remaining players, say t , is even there is an optimal (from the point of view of the best remaining player) format that calls for $t/2$ matches in the next round. We also show, for the section's special case, that the worst format for the best player is to have exactly one match each round. Analogous results for the worst player are also shown. Although we do not have a proof, we conjecture that, among all possible formats, the balanced format maximizes and the one-match-per-round format minimizes the best player's probability of winning the tournament even in the case of general v_i . (We show by a counterexample that the format that calls for $t/2$ matches when an even number t of players remain is not optimal for the best player in the general case.)

2. Literature Review

Most papers in the literature on random knockout tournaments consider more structured formats than the ones we are considering which suppose that the number of matches in each round is fixed

and that the game participants in each round are randomly chosen from those that remain. In these papers the structure of the tournament is fixed and players are randomly assigned to positions of the structure. For example, a structure with 6 players having a balanced format (meaning 2 matches in round 1, 2 matches in round 2 and 1 match in round 3) might in round 1 have a match between those in positions 1 and 2 and one between those in positions 3 and 4, with the winners then playing each other in round 2 and 5 playing 6 in round 2. Another possibility would be to have the winners of round 1 each play in round 2 one of those that did not play in round 1. Figures 1a and 1b indicate these two structures.




Figure 1 Two balanced structures with 6 players

An example of a non-balanced structure with 6 players is one where there is a single match in each round, with the winner of a match playing in the following round someone who has not yet played.

Maurer [10] proved for random structured formats, when $v_1 > v = v_2 = v_3 = \dots = v_n$, that the win probability of Player 1 is maximized under the balanced structures. Because, as noted by Maurer, a random format can be expressed as a mixture of random structured formats, this also establishes the result for random formats. In section 5 we give another proof of this result, which is both quite elementary and also shows, under the same condition, that when there are $n = 2^s + k$ remaining players the optimal random format can schedule either k or $n/2$ games if n is even, and k games

if n is odd. We also prove, when $v_1 > v = v_2 = v_3 = \dots = v_n$, that the win probability of Player 1 is uniquely minimized by the random format that has exactly one match in each round.

Chung and Hwang [1] proved that for any structured format, if v_i is decreasing in i then so is the winning probability, which also proves the result for any random unstructured format. In Corollary 1 of Section 4 we give a proof of this result that also proves, when v_i is decreasing in i , that the probability Player i reaches round s is decreasing in i for all s .

There are other papers studying knockout tournaments, for example, David [2], Glenn [3], Hennessey and Glickman [4], Horen and Riezman [5], Hwang [6], Marchand [9], Khatibi et al. [8], Jacobson et al. [7]. The paper Ross and Ghamami [11] discusses efficient ways of simulating win probabilities for such tournaments.

3. Lower Bound on the Strongest Player's Win Probability

In this section we establish a lower bound on the probability that the strongest player wins the tournament, and an upper bound on the probability that the weakest player wins.

THEOREM 1. *If v_i is decreasing in i , then*

$$P_1 \geq \frac{v_1}{\sum_{j=1}^n v_j}, \quad P_n \leq \frac{v_n}{\sum_{j=1}^n v_j}.$$

Preliminary to proving the theorem, we define the *weight* of a player as follows. Say that a player is alive if that player has not been eliminated. If S is the current set of alive players then, for $i \in S$, the weight of Player i is defined to equal $v_i / \sum_{j \in S} v_j$. If i is no longer alive, its weight is defined to be 0.

The following Lemma will be needed to prove the theorem.

LEMMA 1. *Suppose at the start of a round that players A and B are alive and have respective weights x and y , where $x \geq y$. If X and Y are random variables denoting the respective weights of A and B after the round, then $yE[X] \geq xE[Y]$.*

Proof: The result is immediate if $y = 0$ so suppose that $x \geq y > 0$. Define a random variable Z whose value is determined by the results of the following round, as follows:

- $Z = 0$ if A and B both lose
- $Z = 1$ if A and B both advance
- $Z = 2$ if A and B play each other
- $Z = (i, j)$ if one of the players A or B defeats i whereas the other loses to j
- $Z = (0, j)$ if one of the players A or B advances by not being selected to play whereas the other loses to j

By considering the possible values of Z , we now show that $yE[X|Z] \geq xE[Y|Z]$.

1. $E[X|Z = 0] = E[Y|Z = 0] = 0$

2. Conditional on $Z = 1$, let R represent the sum of the weights of those aside from A and B that advance. Then,

$$E[X|Z = 1] = E\left[\frac{x}{x+y+R}\right], \quad E[Y|Z = 1] = E\left[\frac{y}{x+y+R}\right]$$

showing that $yE[X|Z = 1] = xE[Y|Z = 1]$.

3. Conditional on $Z = 2$, let R represent the sum of the weights of those aside from A and B that advance (and note that R is independent of who wins the game between A and B). Then,

$$E[X|Z = 2] = \frac{x}{x+y} E\left[\frac{x}{x+R}\right], \quad E[Y|Z = 2] = \frac{y}{x+y} E\left[\frac{y}{y+R}\right]$$

Hence,

$$yE[X|Z = 2] - xE[Y|Z = 2] = \frac{xy}{x+y} E\left[\frac{x}{x+R} - \frac{y}{y+R}\right] \geq 0$$

where the inequality follows because $x \geq y$ implies that $\frac{x}{x+r} \geq \frac{y}{y+r}$ for any $r \geq 0$.

4. With w and v representing, respectively, the weights of i and j before the round, and using the fact that A is equally likely to play i or j , we have

$$P(A \text{ advances} | Z = (i, j)) = c \left(\frac{x}{x+w} \right) \left(\frac{v}{v+y} \right)$$

where $1/c = \left(\frac{x}{x+w} \right) \left(\frac{v}{v+y} \right) + \left(\frac{y}{y+w} \right) \left(\frac{v}{v+x} \right)$. Let R represent the sum of the weights of those, aside from A, B and j , that advance, and note that R is independent of which of A or B loses. Then

$$E[X | Z = (i, j)] = c \left(\frac{x}{x+w} \right) \left(\frac{v}{v+y} \right) E \left[\frac{x}{R+x+v} \right]$$

and, similarly,

$$E[Y | Z = (i, j)] = c \left(\frac{y}{y+w} \right) \left(\frac{v}{v+x} \right) E \left[\frac{y}{R+y+v} \right]$$

Hence, letting $D = yE[X | Z = (i, j)] - xE[Y | Z = (i, j)]$, we see that

$$D = cxyv E \left[\frac{x}{(x+w)(y+v)(R+x+v)} - \frac{y}{(y+w)(x+v)(R+y+v)} \right]$$

As it is easy to check that for every $R \geq 0$ and $x \geq y$

$$x(y+w)(x+v)(R+y+v) \geq y(x+w)(y+v)(R+x+v)$$

it follows that $yE[X | Z = (i, j)] - xE[Y | Z = (i, j)] \geq 0$ for $x \geq y$.

5. That $yE[X | Z = (0, j)] - xE[Y | Z = (0, j)] \geq 0$ follows from the preceding result by setting $w = 0$.

Hence, we have shown that $yE[X | Z] \geq xE[Y | Z]$, and the result follows upon taking expectations of both sides of this inequality. ■

Proof of Theorem 1: We give the proof that $P_1 \geq v_1 / \sum_{j=1}^n v_j$. The proof that $P_n \leq v_n / \sum_{j=1}^n v_j$ is similar. Let $W_j(k)$ be the weight of Player j after k rounds have been played. Also, let H_k be the history of all results through the first k rounds. We claim that

$$E[W_1(k+1) | H_k] \geq W_1(k)$$

Because the claim is true when $W_1(k) = 0$, assume that $W_1(k) > 0$. Let A_k denote the set of alive players after round r . Now, from Lemma 1, we have, for $j \in A_k$, that

$$W_j(k)E[W_1(k+1)|H_k] \geq W_1(k)E[W_j(k+1)|H_k]$$

Hence, if $E[W_1(k+1)|H_k] < W_1(k)$, then for any $j \in A_k$

$$W_j(k) > E[W_j(k+1)|H_k]$$

which is a contradiction since $1 = \sum_{j \in A_k} W_j(k) = \sum_{j \in A_k} E[W_j(k+1)|H_k]$. Hence, $E[W_1(k+1)|H_k] \geq W_1(k)$, and taking expectations of both sides gives

$$E[W_1(k+1)] \geq E[W_1(k)], \quad r \geq 0.$$

If the tournament has r rounds, then the preceding yields that $E[W_1(r)] \geq E[W_1(0)]$, which gives the result since $E[W_1(r)] = P_1$ whereas $W_1(0) = v_1 / \sum_{j=1}^n v_j$. ■

Remark: The preceding argument shows that $W_1(k), k \geq 0$ is a submartingale, and that $W_n(k), k \geq 0$ is a supermartingale.

4. Bounds on Win Probabilities

Let

$$p_i = \frac{1}{n-1} \sum_{j \neq i} \frac{v_i}{v_i + v_j}$$

be the probability that i would win a match against a randomly chosen opponent. In this section we prove that P_i is smaller than it would be if it were the case that i would win each game it plays with probability p_i . That is, we will prove the following.

THEOREM 2. *If the tournament format is (r, m_1, \dots, m_r) then*

$$P_i \leq \prod_{s=1}^r \left(\frac{2m_s}{r_s} p_i + 1 - \frac{2m_s}{r_s} \right)$$

where $r_s = n - \sum_{j=1}^{s-1} m_j$ is the number of players that advance to round s .

To prove the preceding theorem we will need a couple of lemmas. Before giving these lemmas, we introduce the following notation. We let $R_{i,s}$ be the event that Player i reaches round s , $s = 1, \dots, r$, and let $R_{i,r+1}$ be the event that i wins the tournament. If a player receives a bye in a round (that is, if it reaches that round but is not chosen to play a match) say that it plays Player 0 in that round. Also, let $p_{ij} = \frac{v_i}{v_i + v_j}$, $i \neq j$, be the probability that i beats j in a game.

Lemma 2 is easily proven by a coupling argument.

LEMMA 2. *For all $s = 1, \dots, r + 1$, $P(R_{i,s}R_{j,s})$ is an increasing function of v_j . (When $i = j$, this states that $P(R_{i,s})$ is an increasing function of v_i .)*

LEMMA 3. *Assume $v_1 \geq v_2 \geq \dots \geq v_n$. For fixed i , $P(R_{i,s}R_{j,s})$ decreases in j for $j \neq i$.*

Proof: Fix i, j, k , where $j < k$ and $j, k \neq i$. The proof is by induction on n . For $n = 3$, the only format is to play one game each round. The result holds in this case because $v_j \geq v_k$ implies that

$$3P(R_{i,2}R_{j,2}) = \frac{v_i}{v_i + v_k} + \frac{v_j}{v_j + v_k} \geq \frac{v_i}{v_i + v_j} + \frac{v_k}{v_j + v_k} = 3P(R_{i,2}R_{k,2})$$

Assume the results holds for up to $n - 1$ players and for all formats. We now consider the n player case. Let π be an arbitrary format. Define a random vector Z whose value is determined by the results of the first round of the tournament under π , as follows:

- $Z = (1, u)$ if i plays against j and k plays against u , or i plays against k and j plays against u ;
- $Z = (2, u)$ if i plays against u , and j plays against k ;
- $Z = (3, u, v, w)$ if i plays against u , j beats v , and k beats w ;
- $Z = (4, u, v, w)$ if i plays against u , j loses to v , and k loses to w ;
- $Z = (5, u, v, w)$ if either (a): i plays against u , j beats v , and k loses to w , or (b): i plays against u , k beats v , and j loses to w .

By considering the possible values of Z , we now show that $P(R_{i,s}R_{j,s}|Z) \geq P(R_{i,s}R_{k,s}|Z)$.

1. Let $p_{zy} = \frac{v_z}{v_z+v_y}$. Because $p_{ik} \geq p_{ij}$ and $p_{ju} \geq p_{ku}$, it follows that

$$P(R_{i,2}R_{j,2}|Z = (1, u)) = p_{ik}p_{ju}/2 \geq p_{ij}p_{ku}/2 = P(R_{i,2}R_{k,2}|Z = (1, u)).$$

Let T denote the set of players other than i, j, k , that reach round 2. Given that i and j have reached round 2, the probability of i and j reaching round $s > 2$ equals to the probability that i and j reaches round $s - 1$ in a new tournament which begins with players i, j and T , and follows the same format as π after round 1. Let $P_{i,j,T}(s)$ denote the probability that i and j reaches round $s - 1$ in the new tournament. Similarly, let $P_{i,k,T}(s - 1)$ denote the probability that i and k reaches round $s - 1$ in a tournament which begins with players i, k and T , and follows the same format as π after round 1. Then by Lemma 2, $P_{i,j,T}(s - 1) \geq P_{i,k,T}(s - 1)$. Therefore for $s > 2$,

$$\begin{aligned} P(R_{i,s}R_{j,s}|Z = (1, u)) &= P(R_{i,2}R_{j,2}|Z = (1, u))E[P_{i,j,T}(s - 1)] \\ &\geq P(R_{i,2}R_{k,2}|Z = (1, u))E[P_{i,k,T}(s - 1)] \\ &= P(R_{i,s}R_{k,s}|Z = (1, u)). \end{aligned}$$

where the preceding used that T is independent of the event $R_{i,s}R_{j,s}$ and of the event $R_{i,s}R_{k,s}$.

2. Since $v_j \geq v_k$, we have that

$$P(R_{i,2}R_{j,2}|Z = (2, u)) = p_{iu}p_{jk} \geq p_{iu}p_{kj} = P(R_{i,2}R_{k,2}|Z = (2, u))$$

Let T denote the set of players other than i, j and k that reaches round 2. With the same definition and argument as when $Z = (1, u)$, it can be shown that for $s > 2$,

$$P_{i,j,T}(s) \geq P_{i,k,T}(s)$$

which implies that

$$P(R_{i,s}R_{j,s}|Z = (2, u)) \geq P(R_{i,s}R_{k,s}|Z = (2, u)).$$

3. In this case,

$$P(R_{i,2}R_{j,2}|Z = (3, u, v, w)) = p_{iu} = P(R_{i,2}R_{k,2}|Z = (3, u, v, w)).$$

Now, given that i, j and k have reached round 2, the induction hypothesis implies that the probability of i and j reaching round s is greater than or equal to the probability of i and k reaching round s for $s > 2$. Therefore for $s > 2$,

$$P(R_{i,s}R_{j,s}|Z = (3, u, v, w)) \geq P(R_{i,s}R_{k,s}|Z = (3, u, v, w)).$$

4. In this case, $P(R_{i,s}R_{j,s}|Z = (4, u, v, w)) = P(R_{i,s}R_{k,s}|Z = (4, u, v, w)) = 0$.

5. In this case,

$$P(R_{i,2}R_{j,2}|Z = (5, u, v, w)) = cp_{iu}p_{jv}p_{wk}$$

$$P(R_{i,2}R_{k,2}|Z = (5, u, v, w)) = cp_{iu}p_{kv}p_{wj}$$

where $1/c = p_{jv}p_{wk} + p_{kv}p_{wj}$.

Since $p_{jv} \geq p_{kv}$ and $p_{wk} \geq p_{wj}$, we have that

$$P(R_{i,2}R_{j,2}|Z = (5, u, v, w)) \geq P(R_{i,2}R_{k,2}|Z = (5, u, v, w))$$

Let T denote the set of players other than i, j and k that reach round 2, and note that T has the same distribution whether (a) or (b) resulted. With the same definition and argument as when $Z = (1, u)$, it can be shown that for $s > 2$,

$$P_{i,j,T}(s) \geq P_{i,k,T}(s)$$

which implies that

$$P(R_{i,s}R_{j,s}|Z = (5, u, v, w)) \geq P(R_{i,s}R_{k,s}|Z = (5, u, v, w)).$$

Hence, we have shown that $P(R_{i,s}R_{j,s}|Z) \geq P(R_{i,s}R_{k,s}|Z)$, and the result follows upon taking expectations of both sides of this inequality. ■

Proof of Theorem 2: Assume that v_j is decreasing in j . Now, given that i reaches round s , the conditional probability that j also reaches round s is

$$P(R_{j,s}|R_{i,s}) = \frac{P(R_{j,s}R_{i,s})}{P(R_{i,s})}$$

Hence, from Lemma 3 it follows that $P(R_{j,s}|R_{i,s})$ is a decreasing function of j , $j \neq i$. Now,

$$P_i(v_1, \dots, v_n) = P(R_{i,2} \cdots R_{i,r+1}) = \prod_{s=1}^r P(R_{i,s+1}|R_{i,s})$$

Let $C_{i,s}$ be the event that i competes in round s (that is, it is the event that i reaches round s and then plays a match in that round). With $Q_j = P(i \text{ plays } j \text{ in round } s | C_{i,s}), j \neq i$, we have that

$$\begin{aligned} Q_j &= P(i \text{ plays } j \text{ in round } s | C_{i,s} R_{j,s}) P(R_{j,s} | C_{i,s}) \\ &= P(i \text{ plays } j \text{ in round } s | C_{i,s} R_{j,s}) P(R_{j,s} | R_{i,s}) \\ &= \frac{1}{r_s - 1} P(R_{j,s} | R_{i,s}) \end{aligned}$$

where $r_s = n - \sum_{j=1}^{s-1} m_j$. Hence, $Q_j, j \neq i$ is a decreasing function of j . Letting Y be a random variable such that $P(Y = j) = Q_j, j \neq i$, it thus follows that Y is stochastically smaller than the random variable X having $P(X = j) = 1/(n-1), j \neq i$. Therefore,

$$\begin{aligned} P(R_{i,s+1}|R_{i,s}) &= 1 - \frac{2m_s}{r_s} + \frac{2m_s}{r_s} P(R_{i,s+1}|C_{i,s}) \\ &= 1 - \frac{2m_s}{r_s} + \frac{2m_s}{r_s} \sum_{j \neq i} \frac{v_i}{v_i + v_j} Q_j \\ &= 1 - \frac{2m_s}{r_s} + \frac{2m_s}{r_s} E\left[\frac{v_i}{v_i + v_Y}\right] \\ &\leq 1 - \frac{2m_s}{r_s} + \frac{2m_s}{r_s} E\left[\frac{v_i}{v_i + v_X}\right] \\ &= 1 - \frac{2m_s}{r_s} + \frac{2m_s}{r_s} \frac{1}{n-1} \sum_{j \neq i} \frac{v_i}{v_i + v_j} \end{aligned}$$

where the preceding used that $\frac{v_i}{v_i + v_j}$ is an increasing function of j and X is stochastically larger than Y to conclude that $E\left[\frac{v_i}{v_i + v_Y}\right] \leq E\left[\frac{v_i}{v_i + v_X}\right]$. ■

Remark: It follows from Lemma 2 that for all random formats, $P_i = P(R_{i,r+1})$ is an increasing function of v_i . On the other hand, it seems intuitive that P_i is a decreasing function of v_j for $j \neq i$. However, while this is true for $n = 3$, it is not true for $n \geq 4$. The argument, when $n = 3$, uses that

$$P_1 = \frac{1}{3} \left(\frac{v_2}{v_2 + v_3} \frac{v_1}{v_1 + v_2} + \frac{v_3}{v_2 + v_3} \frac{v_1}{v_1 + v_3} \right) + \frac{2}{3} \frac{v_1}{v_1 + v_2} \frac{v_1}{v_1 + v_3}.$$

This gives

$$\begin{aligned} \frac{\partial P_1}{\partial v_2} &= \frac{1}{3} v_1 \frac{\partial}{\partial v_2} \frac{v_2}{(v_2 + v_3)(v_1 + v_2)} + \frac{1}{3} \frac{v_1 v_3}{v_1 + v_3} \frac{\partial}{\partial v_2} \frac{1}{v_2 + v_3} + \frac{2}{3} \frac{v_1^2}{v_1 + v_3} \frac{\partial}{\partial v_2} \frac{1}{v_1 + v_2} \\ &= \frac{1}{3} \frac{v_1}{(v_1 + v_2)^2 (v_2 + v_3)^2 (v_1 + v_3)} \left(-v_1 v_3^2 - 2v_2^2 v_3 - 3v_1 v_2^2 - 6v_1 v_2 v_3 \right) \\ &\leq 0. \end{aligned}$$

For $n = 4$ a counter example can be constructed as follows. Consider the balanced format and let $P_i(v_1, v_2, v_3, v_4)$ denote the probability that i wins the tournament when Player j has value v_j , $j = 1, 2, 3, 4$. Conditioning on whether or not Player 1 first plays against Player 4, we have

$$P_1(2, 1, 1, x) = \frac{1}{3} \left(\frac{2}{2+x} \cdot \frac{2}{3} \right) + \frac{2}{3} \cdot \frac{2}{3} \left(\frac{1}{1+x} \cdot \frac{2}{3} + \frac{x}{(1+x)(2+x)} \right)$$

and thus

$$P_1(2, 1, 1, \frac{1}{100}) = \frac{31600}{60903} \approx .518858\dots < .518861\dots \approx \frac{7744}{14925} = P_1(2, 1, 1, \frac{1}{99}). \quad \blacksquare$$

Using Lemma 3, it is easy to show that if v_i is decreasing in i then so is $P(R_{i,s})$.

COROLLARY 1. *If $v_1 \geq \dots \geq v_n$ then $P(R_{1,s}) \geq P(R_{2,s}) \geq \dots \geq P(R_{n,s})$, $s = 1, \dots, r + 1$.*

Proof: Suppose $i < j$. Because r_s is the number of players that are still alive at the start of round s , it follows that given i reaches round s , the expected number of others that also reach round s is $r_s - 1$. Consequently,

$$r_s - 1 = \sum_{k \neq i} P(R_{k,s} | R_{i,s})$$

Hence,

$$(r_s - 1)P(R_{i,s}) = \sum_{k \neq i,j} P(R_{i,s}R_{k,s}) + P(R_{i,s}R_{j,s}),$$

$$(r_s - 1)P(R_{j,s}) = \sum_{k \neq j,i} P(R_{j,s}R_{k,s}) + P(R_{j,s}R_{i,s}),$$

which by Lemma 3 shows the desired result for $s \leq r$. In addition, we have,

$$P(R_{i,r+1}) = \sum_{k \neq j,i} P(R_{i,r}R_{k,r}) \frac{v_i}{v_i + v_k} + P(R_{j,r}R_{i,r}) \frac{v_i}{v_i + v_j},$$

$$P(R_{j,r+1}) = \sum_{k \neq j,i} P(R_{j,r}R_{k,r}) \frac{v_j}{v_j + v_k} + P(R_{i,r}R_{j,r}) \frac{v_j}{v_j + v_i},$$

which by Lemma 3, and by the assumption that $v_i \geq v_j$, completes the proof. ■

We now give lower bounds on the win probabilities.

THEOREM 3. *Suppose $v_1 \geq v_2 \geq \dots \geq v_n$. Let X_1, \dots, X_r be independent with $P(X_j = 1) = \frac{2m_j}{r_j} = 1 - P(X_j = 0)$, $j = 1, \dots, r$, and let $N = \sum_{j=1}^r X_j$. Then*

$$P_i \geq \sum_{g=1}^{i-1} P(N = g) \prod_{k=1}^g \frac{v_i}{v_i + v_k} + \sum_{g=i}^r P(N = g) \prod_{k=1}^{i-1} \frac{v_i}{v_i + v_k} \prod_{k=i+1}^{g+1} \frac{v_i}{v_i + v_k}$$

Proof: Define the surrogate of i as follows: The initial surrogate of i is i itself; anyone who beats a surrogate of i becomes the current surrogate of i . Note that at any time there is exactly one player who is the current surrogate of i . If we let X_j be the indicator of whether the surrogate of i plays a match in round j , it follows that X_1, \dots, X_r are independent with $P(X_j = 1) = \frac{2m_j}{r_j} = 1 - P(X_j = 0)$. Also, let $N = \sum_{j=1}^r X_j$ be the number of games played by surrogates of i (while they are the current surrogate), and let $O = \{J_1, \dots, J_N\}$ be their set of opponents in these games. Because $P(i \text{ wins the tournament} | N, O) = \prod_{k=1}^N \frac{v_i}{v_i + v_{J_k}}$, it follows that

$$P_i = E\left[\prod_{k=1}^N \frac{v_i}{v_i + v_{J_k}}\right]$$

$$\geq E\left[\prod_{k=1}^{\min(N, i-1)} \frac{v_i}{v_i + v_k} \prod_{k=i+1}^{N+1} \frac{v_i}{v_i + v_k}\right]$$

where $\prod_{k=i+1}^{N+1} \frac{v_i}{v_i+v_k}$ is equal to 1 when $N < i$. Hence,

$$P_i \geq \sum_{g=1}^{i-1} P(N=g) \prod_{k=1}^g \frac{v_i}{v_i+v_k} + \sum_{g=i}^r P(N=g) \prod_{k=1}^{i-1} \frac{v_i}{v_i+v_k} \prod_{k=i+1}^{g+1} \frac{v_i}{v_i+v_k} \quad \blacksquare$$

Remarks:

1. The probability mass function of N , which does not depend on i , is easily obtained by solving recursive equations (see Example 3.24 of Ross [12]). In the special case where $n = 2^s + k$, $0 \leq k < 2^s$, and where there are k matches in round 1 and afterwards all remaining players have matches in each subsequent round, we have

$$P(N = s + 1) = \frac{2k}{n} = 1 - P(N = s).$$

2. A weaker bound than the one provided in is obtained by noting that, for $i > 1$, $\frac{v_i}{v_i+v_{J_k}} \geq \frac{v_i}{v_i+v_1}$ and so for $i > 1$

$$\begin{aligned} P_i &\geq E\left[\left(\frac{v_i}{v_i+v_1}\right)^N\right] \\ &= E\left[\left(\frac{v_i}{v_i+v_1}\right)^{X_1+\dots+X_r}\right] \\ &= \prod_{j=1}^r E\left[\left(\frac{v_i}{v_i+v_1}\right)^{X_j}\right] \\ &= \prod_{j=1}^r \left(\frac{2m_j}{r_j} \frac{v_i}{v_i+v_1} + 1 - \frac{2m_j}{r_j}\right) \end{aligned}$$

5. Special Case Best and Worst Formats for the Strongest and Weakest Players

THEOREM 4. *Suppose $v_1 > v = v_2 = v_3 = \dots = v_n$. If $n = 2^s + k$, $0 \leq k < 2^s$, then the balanced format that schedules k matches in round 1 and then has all remaining players competing in each subsequent round leads to the maximal possible value of P_1 .*

Proof: Let $p = \frac{v_1}{v_1+v} > .5$ be the probability that Player 1 wins in a match against another player. The proof is by induction. As there is nothing to prove when $n = 2$, let $n = 2^s + k$, $1 \leq k \leq 2^s$, and suppose that the result is true for all smaller values of n . Consider a format that calls for j matches in the first round, where $j < k$. Then, by the induction hypothesis, the format of this type that is

best for Player 1 will call for $k - j$ matches in the second round. The probability that Player 1 is among the final 2^s players under these conditions is

$$f_1(p) \equiv \left[\frac{2j}{n} p + 1 - \frac{2j}{n} \right] \left[\frac{2(k-j)}{n-j} p + 1 - \frac{2(k-j)}{n-j} \right]$$

However, the probability that Player 1 is among the final 2^s players if the first round has k matches is

$$f_2(p) \equiv \frac{2k}{n} p + 1 - \frac{2k}{n}$$

Now, $f_1(p) - f_2(p) = 0$ when p is either $1/2$ or 1 . Because $f_1(p) - f_2(p)$ is easily seen to be strictly convex this implies that $f_1(p) - f_2(p) < 0$ when $1/2 < p < 1$. Thus, in searching for the best format for Player 1 we need not consider any format that calls for less than k matches in round 1.

Now consider any format that calls for $k + i$ matches in round 1 where $i > 0$. By the induction hypothesis it will call for $2^{s-1} - i$ matches in round 2. This will result in Player 1 being among the final 2^{s-1} players with probability

$$g_1(p) \equiv \left[\frac{2k+2i}{n} p + 1 - \frac{2k+2i}{n} \right] \left[\frac{2^s-2i}{2^s-i} p + 1 - \frac{2^s-2i}{2^s-i} \right]$$

However, the format that calls for k matches in round 1 and 2^{s-1} matches in round 2 leads to Player 1 being among the final 2^{s-1} players with probability

$$g_2(p) \equiv \left(\frac{2k}{n} p + 1 - \frac{2k}{n} \right) p$$

Now, $g_1(p) - g_2(p) = 0$ when p is either $1/2$ or 1 . Because

$$g_1''(p) = 2 \frac{2k+2i}{n} \frac{2^s-2i}{2^s-i}, \quad g_2''(p) = 2 \frac{2k}{n},$$

it follows that $g_1''(p) \geq g_2''(p)$ is equivalent to

$$(k+i)(2^s-2i) \geq k(2^s-i),$$

which is easily seen to be equivalent to

$$2^s \geq k + 2i,$$

which holds because having $k + i$ two person matches implies that $2^s + k \geq 2(i + k)$. Hence, $g_1(p) - g_2(p)$ is convex, which shows that there is an optimal format that initially has k matches in round 1. The induction hypothesis then proves the result. ■

Remarks

1. Because $g_1(p) - g_2(p)$ is strictly convex unless $2^s + k = 2(i + k)$, it follows that the format specified in Theorem 4 is uniquely optimal (in the sense of maximizing P_1) when n is odd, whereas when n is even there is also an optimal format that schedules $n/2$ matches in round 1.

2. In the case of general v_i , the format that calls for $n/2$ matches when an even number n of players remain is not optimal for the best player. For a counter example, consider a knockout tournament with 6 players having values $v_1 = 6, v_2 = 4, v_3 = 3, v_4 = v_5 = v_6 = 1$. Under the balanced format, the probability of Player 1 winning the tournament is $P_1 = 0.4422$. Under a format that plays 3 games in round 1 and then one game in each round, the probability of Player 1 winning the tournament is $P'_1 = 0.4412 < P_1$.

THEOREM 5. *Suppose $v_1 > v = v_2 = v_3 = \dots = v_n$. The unique format that minimizes P_1 is the one that has exactly one match in each round.*

Proof: The proof is by induction. Suppose it is true for all tournaments with fewer than n players and now suppose there are n players. Consider any format that calls for s matches in the first round, where $s > 1$. The probability that Player 1 is still alive when there are only $n - s$ alive players is

$$f_1(p) = \frac{2s}{n}p + 1 - \frac{2s}{n}$$

On the other hand, the probability that Player 1 is still alive when there are only $n - s$ alive players when the format is one match per round is

$$f_2(p) = \prod_{j=0}^{s-1} \left(\frac{2}{n-j}p + 1 - \frac{2}{n-j} \right)$$

Because $f_2(p)$ is a polynomial whose coefficients are all positive it follows that $f_2''(p) > 0$. As $f_1''(p) = 0$ this implies that $f_1(p) - f_2(p)$ is strictly concave, which since $f_1(p) - f_2(p) = 0$ when $p = 1/2$ or $p = 1$, enables us to conclude that $f_1(p) - f_2(p) > 0$ when $1/2 < p < 1$. Hence, by the induction hypothesis, the format that calls for one match in each round results in a win probability for Player 1 that is strictly smaller than what it is under any format that calls for $s > 1$ matches in round 1. ■

The following theorem gives the analogous results for the weakest player. The proofs are similar and thus omitted.

THEOREM 6. *Suppose $v_1 = v_2 = v_3 = \dots = v_{n-1} > v_n$. The format that results in the highest win probability for Player n is the one that has one match in each round. If $n = 2^s + k$, $1 \leq k \leq 2^s$, then the format that minimizes P_n is the one that schedules k matches in round 1 and then has all remaining players competing in each subsequent round.*

We now use the preceding results to obtain universal (that is, they hold for all formats) upper bounds on P_i . Recall that $p_i = \frac{1}{n-1} \sum_{j \neq i} \frac{v_i}{v_i + v_j}$.

LEMMA 4. *Suppose $n = 2^s + k$, $0 \leq k < 2^s$.*

(i) *If $p > \frac{1}{2}$, then $\prod_{j=1}^r \left(\frac{2m_j}{r_j} p + 1 - \frac{2m_j}{r_j} \right)$ is maximized by the balanced format, and its value is $\left(\frac{2k}{n} p + 1 - \frac{2k}{n} \right) p^s$.*

(ii) *If $p = \frac{1}{2}$, then $\prod_{j=1}^r \left(\frac{2m_j}{r_j} p + 1 - \frac{2m_j}{r_j} \right) = \frac{1}{n}$ for all eligible formats.*

(iii) *If $p < \frac{1}{2}$, $\prod_{j=1}^r \left(\frac{2m_j}{r_j} p + 1 - \frac{2m_j}{r_j} \right)$ is maximized by the format where at each round there is exactly one match, and its value is $\prod_{j=1}^{n-1} \left(\frac{2}{n-j+1} p + 1 - \frac{2}{n-j+1} \right)$.*

Proof. Note that for a tournament where the players have respective values (v_1, v, \dots, v) , $P_1 = \prod_{j=1}^r \left(\frac{2m_j}{r_j} p_1 + 1 - \frac{2m_j}{r_j} \right)$. Now, (i) and (iii) are direct corollaries of Theorems 2 and 4 respectively, while (ii) is true because if $v_1 = v$ then all players have equal probability of winning the tournament regardless of the format. ■

THEOREM 7. Suppose $n = 2^s + k$, $0 \leq k < 2^s$.

$$P_i \leq \begin{cases} \left(\frac{2k}{n} p_i + 1 - \frac{2k}{n} \right) p_i^s & \text{if } p_i > \frac{1}{2} \\ \frac{1}{n} & \text{if } p_i = \frac{1}{2} \\ \prod_{j=1}^{n-1} \left(\frac{2}{n-j+1} p_i + 1 - \frac{2}{n-j+1} \right) & \text{if } p_i < \frac{1}{2} \end{cases}$$

Proof. The proof follows directly from Theorem 2 and the preceding Lemma. ■

COROLLARY 2. For a player with $p_i < \frac{1}{2}$, the probability of winning the tournament under any format is less than $\frac{1}{n}$.

Author Biographies

Ilan Adler is a professor of the department of Industrial Engineering and Operations Research at the University of California at Berkeley. Professor Adler hold B.A in Economics and Statistics from the Hebrew University in Israel, M.Sc. in Operations Research from the Technion in Israel and Ph.D in Operations Research from Stanford. His research interests are in optimization theory, financial engineering and combinatorial probability models.

Yang Cao is a PhD student in the Department of Industrial and Systems Engineering at the University of Southern California. He received a Master of Science degree in Industrial and Systems Engineering from University of Southern California, and a Bachelor of Science degree in Theoretical and Applied Mechanics from Peking University.

Richard Karp attended Boston Latin School and Harvard University, receiving the Ph.D. in 1959. From 1959 to 1968 he was a member of the Mathematical Sciences Department at IBM Research. From 1968 to 1994 and from 1999 to the present he has been a professor at UC Berkeley, where he held the Class of 1939 Chair. From 1988 to 1995 and 1999 to the present he has been a Research Scientist at the International Computer Science Institute in Berkeley. His current

activities center on algorithmic methods in genomics and computer networking. He has supervised thirty-six Ph.D. dissertations. Honors and awards include: U.S. National Medal of Science, Turing Award, Fulkerson Prize, Harvey Prize (Technion), Centennial Medal (Harvard), Lanchester Prize, Von Neumann Theory Prize, Von Neumann Lectureship, Distinguished Teaching Award (Berkeley), Faculty Research Lecturer (Berkeley), Miller Research Professor (Berkeley), Babbage Prize and eight honorary degrees. He is a member of the U.S. National Academies of Sciences and Engineering, the American Philosophical Society and the French Academy of Sciences, and a Fellow of the American Academy of Arts and Sciences, the American Association for the Advancement of Science, the Association for Computing Machinery and the Institute for Operations Research and Management Science.

Erol A. Peköz has a BS degree from Cornell University, and MS and Ph.D degrees in Operations Research from University of California, Berkeley. The research of Professor Peköz is focused in three areas: probability approximations and network science, healthcare quality and provider profiling, and queueing models and reliability for operations. Professor Peköz has also taught at Harvard, UCLA and also at UC Berkeley, and is the author of *The Manager's Guide to Statistics*.

Sheldon M. Ross received his B. S. degree in Mathematics from Brooklyn College in 1963. He received his M.S. degrees in Mathematics from Purdue University in 1964 and his Ph.D. degree in Statistics from Stanford University in 1968. Dr. Ross served as a Professor at the University of California, Berkeley from 1976 until joining us in 2004. Dr. Ross also serves as the Editor for Probability in the Engineering and Informational Sciences, the Advisory Editor for International Journal of Quality Technology and Quantitative Management, and an Editorial Board Member of the Journal of Bond Trading and Management. Dr. Ross is the author of Introduction to Mathematical Finance: Options and Other Topics, Simulation, A First Course in Probability, Probability Models for Computer Science, and many more titles, articles and reports.

Acknowledgments

Sheldon M. Ross was supported by, or in part by, the U. S. Army Research Laboratory and the U. S. Army Research Office under contract/grant number W911NF-14-1-0166, and by the National Science Foundation under contract/grant number CMMI1233337.

References

- [1] Chung, F. R. K., and F. K. Hwang, "Do stronger players win more knockout tournaments?," *Journal of the American Statistical Association*, Volume 73, Number 363, 593-596, (1978)
- [2] David, H. A., "Tournaments and paired comparisons," *Biometrika*, Volume 46, Number 1/2, 139-149, (1959)
- [3] Glenn, W. A., "A comparison of the effectiveness of tournaments," *Biometrika*, Volume 47, Number 3/4, 253-262, (1960)
- [4] Hennessy, Jonathan, and Mark Glickman, "Bayesian optimal design of fixed knockout tournament brackets," *Journal of Quantitative Analysis in Sports*, Volume 12, Number 1, 1-15, (2016)
- [5] Horen, Jeff, and Raymond Riezman, "Comparing draws for single elimination tournaments," *Operations Research*, Volume 33, Number 2, 249-262, (1985)
- [6] Hwang, F. K., "New concepts in seeding knockout tournaments," *The American Mathematical Monthly*, Volume 89, Number 4, 235-239, (1982)
- [7] Jacobson, Sheldon H., and Douglas M. King. "Seeding in the NCAA Mens Basketball Tournament: When is a Higher Seed Better?" *Journal of Gambling Business and Economics* 3.2 (2009): 63.
- [8] Khatibi, Arash, Douglas M. King, and Sheldon H. Jacobson. "Modeling the winning seed distribution of the NCAA Division I mens' basketball tournament." *Omega* 50 (2015): 141-148.
- [9] Marchand, Éric, "On the comparison between standard and random knockout tournaments," *Journal of the Royal Statistical Society: Series D (The Statistician)*, Volume 51, Number 2, 169-178, (2002)
- [10] Maurer, Willi, "On most effective tournament plans with fewer games than competitors," *The Annals of Statistics*, 717-727, (1975)
- [11] Ross, S.M., and Samim Ghamami, "Efficient simulation of a random knockout tournament," *Journal Industrial and Systems Engineering*, Volume 2, 88-96, (2008)
- [12] Ross, S.M., *Introduction to Probability Models*, ed. 11, Academic Press, (2014)